

The effect of some organic acid and plant-derived material treatments on the germination and emergence of lettuce

LEVENT ARIN¹, HAYDAR BALCI²

Key words: Lactuca sativa, seed, organic treatments

Abstract

The research was carried out to investigate the effect of plant materials which are used as an alternative to the synthetic chemicals in organic farming (thyme, mint, basil and garlic oil, hot pepper and neem tree seed extract) and some organic acids (salicylic and jasmonic acid) on the germination, emergence and seedling characteristics in lettuce (Lactuca sativa L.) cv Arapsaçı. The seeds of lettuce treated with these materials were subjected to germination and emergence tests at temperature 20±1 °C and 60±5 % RH in autumn period. After that, all seeds (including untreated) were kept at 5±1 °C and again checked for germination, emergence and seedling characteristics in spring period.

Thyme oil affected negatively the germination and emergence and also the infected seed ratio increased. Although the infection ratio was 1% in untreated seeds, it was 7.8% in the seeds treated with thyme oil. After the storing period, infection ratio of thyme oil was 8.5% which was higher than other treatments. There was no significant effect of treatments on seedling characteristics.

Introduction

Lettuce is widely grown both in the field and greenhouse as salad vegetable in all over the world also in Turkey. On the other hand, the size of organically cultivated area in Turkey has reached 614 000 ha and Turkey is one of countries which have highest increases to the area of organic farming all over the world (Willer and Lernoud, 2013).

Because of synthetic-chemical use is forbidden in organic agriculture, some physical and biological methods aimed at the control of seed-borne pathogens are being investigated. There are some physical seed treatment methods like hot water, hot (humid) air, electron beam, but they are required special equipment and have the potential to damage the seed if they are not applied at appropriate dose (Nega et al. 2003, Wolf et al. 2008). The potential alternative seed treatments (microorganism, plant extract and inducer of resistance) have been developing against the different seed-borne pathogens and for mitigating the impact of stress conditions (Groot et al. 2004, Hammer et al. 1999, Klessig and Malamy 1994). However, limited information is currently available dealing with their effect on germination and seedling quality.

The aim of the research was to evaluate the effect of these materials on germination, emergence and seedling quality of lettuce.

Material and methods

The essential oils and plant extracts which were used in this research were selected on basis of their reported antimicrobial properties against a number of pathogens due to be natural agents allowed in organic agriculture. The lettuce seeds had not been treated with any chemicals.

The essential oils were provided from local seller of herbs. The aqueous extracts of hot pepper and neem seed were prepared. The organic acids were obtained from Sigma Aldrich (Interlab Inc. İstanbul). A suspension of essential oils was prepared by dispersing 10 ml amount of essential oil in 1 L of sterile deionized water. The seeds were soaked in water including 10 ml/L oil, extract or acid for 0.5 h at room temperature except jasmonic acid (1 ml/L). After treatment, seeds were dried for one day on a clean cloth under laboratory conditions.

To emergence test, seeds were sown into multi-cell plastic pot (each cell having 30 ml growing media). The properties of peat used as growing medium were as follows: 100-300 ppm N, 100-300 ppm P₂O₅, 150-400 ppm K₂O, pH: 5.4-5.9, electrical conductivity 350 µS/cm. To germination test, each combination of 4 replicates of 100 seed was placed on 2 pieces of Whatman 1 filter paper moistened with distilled water into a 9 cm glass petri dish. Petri dishes were placed in incubator at 20±1 °C, 60±5 % RH with a photoperiod of

¹ Namık Kemal University, Faculty of Agriculture, Turkey, www.nku.edu.tr, eMail: larin@nku.edu.tr

² Yüzüncü Yıl University, Gevaş Vocational School, Turkey, www.yyu.edu.tr, eMail: balcihaydar@hotmail.com

16:8 h (light:dark). Radicle protrusion to 1 mm was scored as germination. Seed germination was recorded at 24 h intervals and germinated seeds were removed from the petri dishes.

The seeds were stored at 5 ± 1 °C for 6 months. Afterwards the experiment was repeated to determine the efficacy of different treatments after storage in spring growing season.

The diagnosis of diseases was made by Prof. Dr. Nuray Özer (Department of Plant Protection, Namık Kemal University) during trials.

The experiment was arranged to completely randomized design with 4 replicates. Arcsin transformation of germination or emergence percentage values was used to stabilize variance. The values were analyzed using analysis of variance followed by an LSD mean separation tests.

Results

There was no statistically significant difference among treatment regarding germination percentage in autumn period, but seed treated thyme oil had the lowest germination percentage in spring and the seeds treated with other materials responded similarly (Table 1). The thyme oil led to the lowest germination percentage with 85.75 %, while other treatments taken part in same important group (ranged from 97.00% to 93.50 %).

In autumn, the mean time germination/emergence was significantly increased after treatment with jasmonic acid compared to other materials (Table 1, 2). It was not observed significant differences among treatments in term of mean time emergence in spring.

The seedling quality properties like seedling length, seedling weight, vigor index etc. was not adversely affected by any of the treatments (data not shown).

The seed treated with thyme oil have highest percentage of infected seed. When it taken into account germination, emergence, seedling properties and infection ratio, best results was obtain from seeds treated with neem tree extract (Table 1, 2).

Table 1: Effect of different seed treatments on germination percentage (GP, %), mean time germination (MTG, days) in autumn and spring periods

Treatments	GP		MTG	
	Autumn	Spring	Autumn	Spring
Thyme oil	92.00	85.75	4.55	3.71
Mint oil	92.00	93.50	4.20	3.67
Basil oil	93.25	94.50	4.13	3.48
Garlic oil	91.75	95.50	4.05	3.49
Hot pepper extract	97.50	93.50	4.10	3.37
Neem seed extract	94.75	94.50	4.08	3.31
Salicylic acid	92.25	94.25	4.01	3.38
Jasmonic acid	94.75	97.00	4.86	3.71
Non-treated	94.50	96.25	4.11	3.47
		*	**	**

* significant at $P<0.05$ and ** significant at $P<0.01$

Table 2: Effect of different seed treatments on emergence percentage (EP, %), mean time emergence (MTE, days) and infected seedling ratio (ISR, %) in autumn and spring periods

Treatments	EP		MTE		ISR	
	Autumn	Spring	Autumn	Spring	Autumn	Spring
Thyme oil	84.00	85.75	7.69	9.94	7.75	8.50
Mint oil	71.00	80.50	7.43	8.99	2.00	3.50
Basil oil	79.00	90.25	7.40	8.60	1.00	3.25
Garlic oil	81.00	87.50	7.37	8.32	1.33	2.25
Hot pepper extract	90.00	84.50	7.07	9.11	1.50	4.25
Neem seed extract	95.00	88.75	7.09	7.87	1.00	2.75
Salicylic acid	82.50	90.00	7.57	8.40	2.00	2.75
Jasmonic acid	75.50	87.50	8.17	9.04	1.66	4.00
Non-treated	76.50	90.25	7.69	8.32	1.00	2.00
			*		**	*

* significant at $P < 0.05$ and ** significant at $P < 0.01$

Discussion

As similar to results concerning mean germination time, Blum et al. (2006) stated that there was no indication of a positive effect of seed treatment with plant strengtheners on parsley or dill concerning a better germination rate of emergence

Although thyme oil, among essential oils tested, exhibited the highest in vitro inhibitory activity against seed-borne pathogens (Wolf et al. 2008), in this research, the seed treated with thyme oil have highest percentage of infected seed, while infection ratio changed between 1% and 4% in others. This may be due to high concentration and/or the lack of sufficient purity. Wolf et al. (2008) reported that cinnamon oil at a concentration of 3.3% was damaged the cabbage seeds.

Although neem is generally known to be effective against insects, it was reported that neem can be used for Fusarium wilt by Foerster et al. (2001).

Suggestions

Based on these results, it can be stated that the neem seed extract can be used to promote germination or emergence and to decrease disease infection without loss of quality characteristics of seedlings.

References

- Blum H, Fausten G, Nega E, Jahn M, Garber U & Aedtner L (2006): Improvement of seed quality of medicinal plants and herbs in organic farming. The Joint European Organic Congress, May 30-31 2006, Odense, Denmark.
- Foerster P, Varela A & Roth J (2001): Best practices for the introduction of non-synthetic pesticides in selected cropping systems. Deutsche Gesellschaft für Technische Zusammenarbeit GmbH, Eschborn, Germany.
- Groot SPC, Wolf JM van der, Jalink H, Langerak CJ & Bulk R (2004): Challenges for the production of high quality organic seeds. Seed Testing International no.127: pp 12-15.
- Hammer KA, Carson CF & Riley TV (1999): Antimicrobial activity of essential oils and other plant extracts. *Journal of Applied Microbiology*, 86, 985-990.
- Klessig DF & Malamy J (1994): The salicylic acid signal in plants. *Plant Molecular Biology*, 26(5): 1439-1458.
- Nega E, Ulrich R, Werner S & Jahn M (2003): Hot water treatment of vegetable seed – an alternative seed treatment method to control seed-borne pathogens in organic farming. *Journal of Plant Diseases and Protection*, 110 (3): 220-234.
- Willer H & Lernoud J (2013): *Organic Agriculture Worldwide: Key results from the FiBL-IFOAM survey on organic agriculture worldwide 2013* (www.fibl.org).
- Wolf JM van der, Birnbaum YE, Zouwen PS van der & Groot SPC (2008): Disinfection of vegetable seed by treatment with essential oils, organic acids and plant extract. *Seed Science and Technology*, 36, 76-88.

