

It was then assumed that the background N_2O emission as well as the N_2/N_2O -ratios as a function of clay content could be described with a Michaelis-Menten equation (Fig. 6). The background N_2O emission was fitted with this equation to give an average emission at about $1 \text{ kg N ha}^{-1} \text{ year}^{-1}$ (Fig. 1). Similarly, the N_2/N_2O ratios were fitted resulting in an average N_2/N_2O ratio at about 4, which corresponds to the mean value of the about 500 N_2/N_2O -ratios found in the literature.




Fig. 2. Relations between water filled pore space (WFPS) and N_2/N_2O ratio in four different soils (from Weier *et al.*, 1993).




Fig. 3. Relations between water filled pore space (WFPS) and denitrification activity.




Fig. 4. Relationship between clay content and WFPS at field capacity.




Fig. 5. Relationship between clay content and saturated hydraulic conductivity.

SimDen is described in further details in (Vinther & Hansen, 2004) and can be downloaded at www.agrsci.dk/simden

Results

SimDen was compared with a number of field measurements in Danish soils (Fig. 7).




Fig. 6. Fitted relationships between background N_2O emission (a) and N_2/N_2O -ratios (b) as function of clay content. Dots are measured values of WFPS at field capacity as function of clay content.

Within the entire range of measurements (Fig. 7, left) there seem to be a reasonable good agreement between the measured denitrification rates and those calculated with SimDen. The high denitrification rate at $230\ kg\ N\ ha^{-1}\ year^{-1}$ was measured during the third year in a loamy soil (ST6), which had received $100\ tons\ pig\ slurry\ ha^{-1}\ year^{-1}$ for three years (Maag, 1989). Within the lower range of values, where the major number of results is found (Fig. 7, right), SimDen seems to overestimate the denitrification.

However, different circumstances under which these measurements were made indicate that the measured values may be underestimated: 1) SimDen estimates the denitrification in 0-100 cm, whereas all measurements only include the 0-20 cm soil layer, 2) the measurements in the field were performed with relatively large time intervals, and thus the denitrification flushes following precipitation

may not have been caught, 3) few of the results include measurements during the winter time, where it has been found that up to 75% of the annual emission may be measured (Goossens et al., 2001), and finally, 4) all measurements were performed on well-drained soils without depressions in the field with higher soil moisture and potentials for higher denitrification.




Fig. 7. Measured and SimDen-modelled denitrification rates in the entire range (left) and the lower range of values (right).

Comparisons between the original version of SimDen and SimDen-Clay show a good agreement between the two models (Fig. 8).

Conclusion

Based on knowledge about only clay content and input of fertiliser N is it possible to give a rough estimate of annual denitrification.

If more detailed information is needed, i.e., regarding temporal variation under certain climatic conditions, dynamic models as for example the soil-plant-atmosphere model Daisy must be used.




Fig. 8. Comparisons of average annual denitrification estimated by SimDen and SimDen-Clay at the soil types (ST) 1 – 9 with clay contents from <5% to >45%.

References

- Bouwman, A. F. (1996) Direct emission of nitrous oxide from agricultural soils. *Nutrient Cycling in Agroecosystems* 46, 53-70.
- Goossens, A., De Visscher, A., Boeckx, P. & van Cleemput, O. (2001) Two-year field study on the emission of N₂O from coarse and middle-textured Belgian soils with different land use. *Nutrient Cycling in Agroecosystems* 60, 23-34.
- IPCC (1997) Greenhouse Gas Inventories. Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. *Intergovernmental Panel on Climate Change*.
- Kasimir-Klemetsson, Å. & Klemetsson, L. (2002) A critical analysis of nitrous oxide emissions from animal manure. In: (ed. Petersen S. O. & Olesen J. E.) *Greenhouse gas inventories for agriculture in the Nordic countries. DIAS report 81* Danish Institute of Agricultural Sciences, Foulum, Denmark, pp. 107-121.
- Maag, M. (1989) Denitrification losses from soil receiving pig slurry or fertilizer. In: (ed. Hansen J. Aa. & Henriksen K.) *Nitrogen in Organic Wastes Applied to Soils*. Academic Press, London, pp. 235-246.
- Vinther, F. P. & Hansen, S. (2004) SimDen - a simple empirical model for quantification of N₂O emission and denitrification. *DIAS Report 104*, 1-47 (in Danish with English summary).
- Weier, K. L., Doran, J. W., Power, J. F. & Walters, D. T. (1993) Denitrification and the dinitrogen/nitrous oxide ratio as affected by soil water, available carbon, and nitrate. *Soil Science Society of America Journal* 57, 66-72.