

Proceedings of the



2018

XX INTERNATIONAL

**WORKSHOP
ON SMUTS
AND BUNTS**

Logan, Utah, USA



Bayer CropScience



Ardent Mills®



IDAHO
WHEAT

29 May 2018, Tuesday
Arrival of participants to Logan

Hampton Inn by Hilton
1665 North Main Street
Logan, Utah 84341 USA
Phone +1-435-713-4567

18:30 – 20:30 Welcome Reception:
The Italian Place: 48 Federal Ave, Logan, UT 84321

30 May 2018, Wednesday

09:00–09:30 Welcome to the conference by Dr. Paul Johnson, Department
Head, Plants, Soils and Climate, Utah State University.

09:25–10:00 Introduction and logistics, Utah Bunt nursery details – David
Hole

10:00–10:30 Break.

Scientific session:
Pathogens: Diversity, Pathogenicity, Methodology
Session chair: Hermann Burstmayr

10:30–11:00 **iTRAQ-based Proteomic Analysis of Wheat Bunt Fungi *Tilletia***
controversa*, *T. caries* and *T. foetida
Li Gao

11:00–11:30 **Classification of wheat bunt diseases (*Tilletia* spp.) and the**
importance of reliable reference material for the development
of new detection methods
Monika K. Grundler

11:30-12:00 **Tracing *Tilletia caries* in wheat during the endophytic phase**
Fabio Mascher

Lunch – 12:00 – 13:30
Session chair: Juliet Marshall

13:30-14:00 **Three new species of flag smut of grasses from the United States**
Kyryll G. Savchenko

14:00-14:30 **Determination of the genome composition of *Sporisorium reilianum* f. sp. *reilianum* (Kühn) Langdon and Fullerton, the sorghum head smut pathogen**
Chunlai Zhang

14:30-15:00 **Historical records of *Urocystis* in North America from the U.S. National Fungus Collections**
Lisa A. Castlebury

15:00-15:30 Break

Sightseeing in Logan or hiking in Logan canyon to wind caves
16:00-

Dinner on your own

31 May 2018, Thursday

**Scientific session:
Disease Control
Session chair: Anders Borgen**

- | | |
|-------------|--|
| 09:00-09:30 | Genome-Wide Association Mapping for Dwarf Bunt Resistance in the National Small Grains Collection
Tyler Gordon |
| 09:30-10:00 | Virulence pattern of Czech bunt samples and sources of resistance
Veronika Dumalasová |
| 10:00-10:30 | Evaluation of Nordic heritage varieties and NILs for resistance to common bunt (<i>Tilletia caries</i> syn. <i>T. tritici</i>)
Anders Borgen |

10:30-11:00 Break
Session chair: Veronika Dumalasova

- | | |
|-------------|---|
| 11:00-11:30 | Strategic use of virulence pattern to develop genetic markers for resistance to common bunt (<i>Tilletia caries</i>) in wheat
Anders Borgen |
| 11:30-12:00 | New tools available to control the common bunt of wheat: development of an early detection test on plantlet by qPCR.
G. Orgeur. |
| 12:00-12:30 | Mapping QTLs conferring additive resistance to Karnal bunt in bread wheat in two recombinant inbred lines populations
P.K. Singh |

12:30 – 14:00 – Lunch
Session chair: Fabio Mascher

14:00-14:30 **Prospects and challenges for breeding bunt resistant wheat
using molecular marker assisted selection**
Rui Wang

14:30-15:00 **Comparative mapping of bunt resistance genes in winter wheat**
Hermann Bürstmayr

15:00-15:30 **Organic methods of controlling common bunt at the farm level.**
Lars Wiik

15:30-16:00 Break
Session chair: David Hole

16:00-17:00 Business Meeting - Discussion

Banquet – 19:00 – 21:00
Smithfield Golf Course Reception

01 June 2018, Friday

09:00-09:30	Travel to Logan Bunt nursery
10:30-12:00	Travel to Pocatello, ID
12:00-13:00	Lunch on own at Portneuf Valley Brewing
13:30-16:00	USDA small grains collection Aberdeen, ID (Passport or Gov't ID required)
16:00	Travel back to Logan or SLC airport hotels.

Strategic use of virulence pattern to develop genetic markers for resistance to common bunt (*Tilletia caries*) in wheat

Anders Borgen¹, Gunter Backes², Karl-Josef Müller³ and Hartmut Spieß⁴

¹ *Agrologica, Houvej 55, DK-9550, Denmark*

² *University of Kassel, FB 11 - Organic Agricultural Sciences, Steinstr. 19, D-37213 Witzenhausen, Germany*

³ *Getreidezüchtungsforschung Darzau, Hof Darzau 1, D-29490 Neu Darchau, Germany*

⁴ *Landbauschule Dottenfelderhof e.V., Dottenfelder Hof 1, 61118 Bad Vilbel, Germany*

borgen@agrologica.dk

When assessing races of common bunt for virulence pattern within a region, it is important to take into account that collected spores may represent a diverse population of different virulence races. When screening spores on a differential set of wheat lines with known resistance genes, a low infection rate on a resistant wheat variety does not necessarily demonstrate that virulence is absent in the spore collection, but could be a sign that virulence is present, but only present in a low frequency among the spores. If just a few spores within a spore sample are indeed virulent, they may infect some plants and from there multiply the virulence quite rapidly next years. Previous studies have shown that virulence against most resistance genes were present in Denmark after purifying races of common bunt (*Tilletia caries*) on resistant varieties. So far, only wheat differential varieties with *Bt4*, *Bt6*, *Bt9*, *Bt11* and *Bt12* cannot be infected with bunt races purified from Danish collections [1, and later own unpublished data]. Virulence against *Bt4*, *Bt6* and *Bt9* has been found in other European studies [2], and *Bt11* may not be only one gene but a combination of at least two genes [3]. Therefore, *Bt12* seems to be the only gene for which virulence have not been found in European population of common bunt. This leads to the conclusion that if resistance breeding shall safely control common bunt in wheat, we need not only one effective gene, but a combination of pyramided genes. Since it is very difficult to test if a resistant line has only one gene or more genes, the most effective tool to achieve this at present are genetic markers.

Using Genome Wide Association Studies (GWAS) to find QTLs and markers for the major resistance genes in wheat have so far led to only few commercial useful markers. Till now, only markers for *Bt9* [6] and *Bt10* are used in practice, but a marker for *Bt12* [4] and Blizzard [7] have also been found. One of the problems in developing markers for bunt resistance have been that spores used in GWAS trials have been diverse or unknown in virulence, and that phenotypic results not distinguishes between different resistance genes. Therefore, the most successful studies have used segregating populations of single crosses where the resistance gene is known on before hand [5].

In the LIVESEED project, we have the ambition to develop genetic markers on several different resistance genes at the same time. We will do so by testing segregating populations of several different crosses between varieties with 7 different resistance genes, and infect them with 7-11 different virulence races of common bunt able to distinguish between the resistance genes. A total of 300 varieties will be pheno- and genotyped. Using this experimental design, we attempt during

2018 and '19 to develop markers for *Bt1*, *Bt2*, *Bt5*, *Bt7*, *Bt13*, *BtZ* and Quebon-resistance, and hopefully also a couple of minor QTLs.

Acknowledgement: The research is part of the LIVESEED project supported by EU Horizon2020 program.

References

- [1] Borgen, A. 2016. Screening wheat varieties for resistance with purified virulence races of common bunt (*Tilletia caries*). Abstract of the XIX international Workshop on Smuts and bunts. Ed.: Alexey Morgounov, Hafiz Muminjanov P 27-30
- [2] Dumasová, V and P. Bartoš 2013. Wheat screening for resistance to common bunt and dwarf bunt. 63. Tagung der Vereinigung der Pflanzenzüchter und Saatgutkaufleute Österreichs 2012. p51-54
- [3] Hagenguth, J.F. 2016: Breeding for organic agriculture: Evaluation of common bunt resistance in three winter wheat populations derived from M822123, PI178383 and PI560841-bcl. MSc-Thesis, BOKU, Austria p.145.
- [4] Müllner, A.E, B. Eshonkulov, J. Hagenguth, B. Pachler, H. Huss, D. Hole, H. Buerstmayr 2016. Comparative mapping of common bunt and dwarf bunt resistance QTL in winter wheat. 67. Tagung der Vereinigung der Pflanzenzüchter und Saatgutkaufleute Österreichs p.39
- [5] Rasmussen, S.K., P. M. Steffan, A.M.Torp, A. Borgen, G.Backes 2016. Mapping resistance genes for common bunt in wheat. Abstract of the XIX international Workshop on Smuts and bunts. Ed.: Alexey Morgounov, Hafiz Muminjanov p: 6-10
- [6] Steffan, P., A.M. Torp, A.Borgen, G.Backes, S.K. Rasmussen 2017: Mapping of common bunt resistance gene *Bt9* in wheat. Theoretical and Applied Genetics. 130(5,):1031–1040
- [7] Wang S1, Knox RE, DePauw RM, Clarke FR, Clarke JM, Thomas JB., 2009: Markers to a common bunt resistance gene derived from 'Blizzard' wheat (*Triticum aestivum* L.) and mapped to chromosome arm 1BS. Theoretical and Applied Genetics 119(3):541-53.