

# Bioactive forage and phytotherapy to cure and control endo-parasite diseases in sheep and goat farming systems – a review of current scientific knowledge

Gerold Rahmann and Hannah Seip

Institute of Organic Farming, Agricultural Research Centre (FAL), D-23847 Trenthorst, [oel@fal.de](mailto:oel@fal.de)

## Abstract

Infestation with gastro-intestinal nematodes (GIN) in small ruminants can cause server economic losses and endanger animal welfare. The development of organic farming systems, the increased public awareness for drug residues in agricultural products and the development of resistant strains of parasites have enforced the search for sustainable alternatives.

The aim of this review is to summarise the current scientific knowledge of alternative strategies to prevent and control endo-parasitic diseases in organic sheep and goat farming systems. The conducted literature evaluation has revealed the major potential to be within the field of bioactive forages, phytotherapy, homeopathy and copper-oxide wire particles. Alternative management pattern like grazing management, nematophagous fungi, improved fodder and breeding are not considered.

The administration and cultivation of bioactive forages and phytotherapy are displaying promise potential for endo-parasite control in organic sheep and goat farming. Scientific research has mainly concentrated on the extracts of the plant species chicory, Birdsfoot trefoil (*Lotus corniculatus*), Sainfoin (*Onobrychis viciifolia*), Sulla (*Hedysarum coronarium*) and Quebracho (*Schinopsis ssp.*). The analysis of these plants showed all plants to have some positive potential, but also highlighted individual limitations in application. However from the results of this literature review none of the investigated plants have been researched sufficiently in on farm experiments to recommend any for implementation at this stage.

No concrete recommendation for a single plant can be given, further research on promising species for the commercial use is strongly recommended, as is the review of the law concerning the appliance of plant based remedies.

Key words: endo-parasite, bioactive forage, phytotherapy, organic farming, small ruminants

## **Stand des Wissens über bioaktives Futter und Phytotherapie zur Behandlung und Kontrolle von Endoparasitosen in der Schaf- und Ziegenhaltung.**

Infektionen mit Endoparasiten – insbesondere den Magen-Darm-Strongyliden (MDS) - sind eine der bedeutendsten Probleme in der Schaf- und Ziegenhaltung. Ziel dieser Literaturstudie war es, den Stand des Wissens über alternative Methoden zur Reduzierung bzw. Kontrolle von MDS festzustellen. Dabei wurden besonders bioaktives Futter, Phytotherapie, Homöopathie und Kupferoxide betrachtet. Alternative präventive Managementmethoden wie optimiertes Weidemanagement, nemathophage Pilzsporen, verbessertes Futter und Zucht wurden hier nicht betrachtet.

Die Anpflanzung von bioaktivem Futter in die Grünlandnarbe (Weide, Wiese) und die Phytotherapie zeigen ein Potenzial für die Kontrolle von Endoparasiten in der Schaf- und Ziegenhaltung. Wissenschaftliche Arbeiten haben sich besonders mit Gewöhnlichem Hornklee (*Lotus corniculatus*), Esparsette (*Onobrychis viciifolia*), Alpen-Süßklee (*Hedysarum coronarium*) und Quebracho (*Schinopsis ssp.*) – einer südamerikanischen Baumart – befasst. Alle zeigten eine gewisse Wirkung, die jedoch nicht überwältigend war. Auch für diese am intensivsten untersuchten Pflanzen sind bislang keine Praxisversuche durchgeführt worden und können damit nicht empfohlen werden. Auch für andere Pflanzen kann bislang keine Empfehlung ausgesprochen werden. Hier fehlen weitergehende Studien, die phytotherapeutische Wirkung belegen. Das Problem besteht auch in der Deklaration, ob es sich um einen Futterzusatzstoff oder ein Medikament handelt. Heilstoffe auf Ganzpflanzen oder Pflanzenteile werden in der Regel nicht mehr als Medikament anerkannt.

Schlüsselwörter: Endoparasitosen, bioaktives Futter, Phytotherapie, Ökologischer Landbau, Kleine Wiederkäuer

## 1 Introduction

Parasitic nematodes remain a major threat to the health and welfare of small ruminants all over the world and the demand for alternative control measures has constantly increased during the last years. Infestation with endo-parasites can have severe consequences for the animal as well as for the livestock farmers leading to economic loss and restricted productivity. Increased public awareness of drug residues in animal products, the increased resistance of parasites to modern anthelmintics, combined with the wish for a more sustainable way of farming has resulted in an intensified effort to find alternative endo-parasite control options (Rahmann et al., 2002). This has triggered the evaluation of some of these traditionally applied plants for their anthelmintic properties with a view to finding out more about them.

Nevertheless, from the scientific point view little has been done in this area, even though a lot of plants are currently used in third world countries (Hördegen, 2005). They are not only dealing with helminth infections but also have to be able to treat a whole range of other diseases with often successful outcomes (Nfi et al., 2001).

The problem is that science has to deal with at the huge variety of plants that may or may not be suitable for the development of alternative anthelmintics. There is also the current lack of verified information to contend with. For these reasons effective plants need to be divided from ineffective ones and the quickest means of achieving this is with in vitro methods. Once a plant has proven its efficiency in vitro, further in vivo testing will be necessary to confirm the obtained results and evaluate risks, side-effects and future applicability (Hounzangebe-Adote et al., 2005a). At this stage it seems a long way off from the discovery of a potential plant to the release of a commercially viable product for use on farms.

The literature review will help researchers, adviser and farmers to get an overview about the state of the art from scientific point of view. Particularly organic farmers are in need to get information about possibilities and limitations of phytotherapy for endo-parasite control.

## 2 Bioactive forages

Plants contain more than energy and protein for animal nutrition but possibly also anti-parasitic compounds, or nutraceuticals. Both terms refer to crops that contain secondary plant substances (SPS) and metabolites which are considered to be beneficial for the animal health rather than having an optimized nutritional value (Rahmann, 2004).

In this context a certain group of SPS, the condensed tannins (CT) have been investigated. CT are not only included in certain plants, a lot of plants have CT content but only those with higher levels are referred to as 'bioactive forage'. As opposed to the application of medical plants, these forages are generally non-toxic and can consequently not be overdosed and the idea is to integrate them into the normal diet of ruminants (Thamsborg, 2001a).

There seem to be several options for feeding tanniferous plants. An example of this is the cultivation of arable crops that can be integrated in the normal rotation (Niezen et al., 1998), these can then be used for either de-worming paddocks, or the plants can be preserved and fed as hay or silage at a later date.

Feeding bioactive forages is not only associated with positive effects but also with some negative consequences (Coop and Kyriazakis, 2001) and this may be the reason why plants with high tanniferous content have not been used earlier. High concentration of CT is known to lead to reduced feed digestibility, feed intake and consequently lower productivity (Aerts et al., 1999; Dawson et al., 1999). Coop and Kyriazakis (2001) therefore conclude that the intake of tanniferous plants will only be preferred by grazing small ruminants "if the negative consequences are offset by the positive effects attributed to their anti-parasitic properties".

Many studies have been undertaken to find out more about the anthelmintic effects of condensed tannins, with two main explanations for the mode of action of condensed tannins being observed:

- First theory: Indirect mode of action: When tannin-rich forages are consumed, the then released condensed tannins build complexes with proteins and protect these from ruminal degradation (tannins have a higher affinity to proteins than to other substances). These complexes dissociate in the abomasum and release protein, ready for absorption. Since nematode parasitism leads to a loss of protein and decreased protein absorption, the intake of tanniferous forages may balance the protein loss and thereby increase resilience. (Min et al., 2003)
- Second Theory: Direct mode of action: Condensed tannins directly react with the proteins on the surface of the parasites and disturb the normal physiological functions of the nematodes like mobility, food absorption or reproduction (Heckendorn, 2005)

For the successful performance a certain CT-content ( $35 \text{ g d}^{-1}$ ) of the applied plants seems to be necessary (Athanasidou et al., 2005). Alternatively Min and Hart (2003) suggest that beneficial effects of CT in plants only occur within the concentration range of 45 to 55 g of CT  $\text{kg}^{-1}$  of DryMatter (DM), levels below and above this range lead to inconsistency.

There is evidence that results obtained in studies with sheep can be transferred to the application on goats (Paolini et al. 2003c; Paolini et al. 2005b). Hoste et al. (2005b) state that data obtained in goat experiments is in agreement with those in sheep. Basic physiological processes are alike indeed but it should be kept in mind that goats are more susceptible than sheep and they take longer to acquire immunity (Thamsborg et al, 2004).

A number of plants have been investigated in the last years and it was found that the CT-content of most grasses is  $<1\%$ , (in DM) most temperate forage legumes have about 5% and some tropical plants contain up to 40% (Thamsborg, 2001a).

## **2.1 Chicory (*Chicorium intybus*, family: *Asteraceae*)**

Chicory is a bushy perennial herb that has light blue to lavender coloured flowers, which is originated in the Mediterranean climate. Forage chicory is known to improve the live weight gain in lambs and to lead to lower pasture contamination because “larval survival on chicory is lower than on grasses” (Ratray, 2003). Similar results were achieved by Marley et al. (2003a, 2003b), who found that infected lambs grazing chicory had the highest live weight gain and the scanned faeces had a tendency of fewer larval development.

Anthanasiadou et al. (2005) proved strong anthelmintic activity in their in vitro studies. In this study extract from rumen material of sheep that previously grazed pure stands of chicory was analyzed, former studies had tested extracts of plant material. The subsequent in vivo study on chicory showed that feeding chicory was indeed effective against adult *T. circumcineta* but not on incoming larvae.

These results were confirmed by a five week grazing trial (Marley et al., 2003a) and two recent studies that investigated that short-term of grazing chicory leads to a reduced adult worm burden but no differences in egg output (Anthanasiadou et al., 2005; Tzamaloukas et al., 2005).

These experiments provide evidence for the ability of chicory to reduce the adult worm burden in infected sheep but they also prove the incapacity of chicory to reduce faecal egg output and prevent incoming larvae from settling. Consequently grazing chicory may help infected animals to balance weight loss due to parasitic infestation and thereby enhance the buildup of immunity. It certainly neither appears suitable to decrease pasture infectivity nor does it show the ability to protect sheep from further infestation.

Unfortunately no reports were found regarding the effectiveness of chicory on goats, so no secured conclusion can be drawn and no recommendation given in this area.

## **2.2 Birdsfoot Trefoil (*Lotus corniculatus*, *L. pendunculatus*, family: *Fabaceae*)**

These two lotus species are herbaceous perennial legumes with yellow flowers that are native to Europe and parts of Asia (Lolicato, 1998). *Lotus pendunculatus* is referred to as ‘Greater Birdsfoot Trefoil’, ‘Maku Lotus’ or ‘Greater Lotus’ whereas the other species is either called ‘Birdsfoot Trefoil’ or ‘Goldie Lotus’.

Ratray (2003) reviewed quite a few of studies on lotus that had been conducted up until 2001. Positive results were found for performance and live weight gain during infections and variable results for the reduction of egg output and worm burdens.

The results of Marley et al. (2003a; 2003b) lead to opposing conclusions. In the feeding trial (Marley et al., 2003a) with lambs carrying mixed infections, they found that the lotus fed group had the lowest weight gain and overall showed no significant positive effect. The impact of different forages on development and survival were explored in their other study (Marley et al., 2003b) and resulted in no effect on hatchability but the highest larval development (L<sub>3</sub>) on birdsfoot trefoil. Two short-term grazing studies had similar results were no direct effects of birdsfoot trefoil could be investigated, although both authors wondered whether the lack of effect might have been due to a minor dosage (Anthanasiadou et al., 2005; Tzamaloukas et al., 2005).

Greater birdsfoot trefoil has also been investigated by Athanasiadou et al. (2005b) within the scope of the wormcops project but the in vitro experiments could not demonstrate any anthelmintic properties. But the in vivo trials measured a substantially declined egg excretion for *T. circumcineta* and *Trichostrongylus spp.*

These results were less clear than those for chicory. Marley et al. (2003a) suggest that inconsistent findings could be related to the fact that different cultivars have varying CT contents. Athanasiadou et al. (2005) and Tzamaloukas et al. (2005) both wonder whether their dosage had been insufficient. Unfortunately no other study has been conducted with different cultivars and levels of CT, to either confirm, or disprove this explanation.

Judging the current stage of experiments, no recommendation for the application in practice can be given; results are inconsistent and controversial and indicate an exploratory need to finally determine the potential of birdfoot trefoil species.

### **2.3 Sulla (*Hedysarum coronarium*, family: Fabaceae)**

Sulla is a biennial or short-lived perennial with flowers that can vary from pink to violet originating from the Western Mediterranean and Northern Africa (Frame, 2006; Frame et al., 1998).

Niezen et al. (1998) found in their six week grazing trial that lambs that had grazed Sulla, had a lower FEC, reduced worm burdens and intestinal parasite density, had good live weight gain, despite infection, and were less affected by parasites.

Molan et al. (2000a) tested the effects of Sulla extract on larval development and found that the genus *T. colubriformis* was more resistant to the inhibitory effects than the other tested species. Further in vitro tests were undertaken within the scope of the WORMCOPS project by Hoste et al. (2005b) who observed reduced in vitro egg hatchability but no difference in larval development. The subsequent in vivo study on lambs infected with *H. contortus* grazing Sulla showed reduced FEC and worm burdens.

Similar results were obtained by Niezen et al. (2002) who discovered that the consumption of Sulla reduced faecal egg output substantially but it also accumulated further evidence for the inefficiency of Sulla on *T. colubriformis*.

The short-term feeding trials of Athanasiadou et al. (2005), Tzamaloukas et al. (2005) and Pomroy and Adlington (2006) do not support any of the previous evidence, both trials show no positive results for Sulla consumption. The former two studies both argue that the lack of effect could be due to insufficient levels of CT, Pomroy and Adlington state similar level of CT in their diet like Niezen et al. (2002) and conclude that the lack of effect must have another reason.

The presented research results indicate that there is a discrepancy between longer and short-term experiments. The administration of Sulla seems to have no effect when applied for a short period of time but if fed for longer, it may reduce FEC and have a positive effect on animal performance.

A further question that should be considered is if it is possible to integrate the cultivation of Sulla into the farming routine. Pomroy and Adlington (2006) state that “Sulla is a difficult herbage to manage agronomically” and they further quote that the provision of a substantial amount of Sulla for an extended period of time for all animals on a farm seems difficult to achieve. Information about Sulla in the FAO-database indicates that weeds can affect Sulla at establishment, as well as after cutting, and that it is not suitable for intensive grazing (Frame, 2006; Frame et al., 1998).

These aspects need to be considered and indication is given that further research is necessary to find out more about the cultivation attributes of Sulla, particularly for organic agricultural systems. Moreover it seems advisable to establish the period of time required for the successful reduction of FEC.

## **2.4 Sainfoin (*Onobrychis viciifolia*, family: *Fabaceae*)**

Sainfoin is a perennial herb which is distributed in Europe, parts of Asia and Northern America (Frame et al., 1998; Frame, 2006).

Barreau et al. (2005) showed in their in vitro experiments the activity of Sainfoin extract against *H. contortus* nematodes and therefore confirmed the results of Paolini et al. (2004) who had demonstrated the inhibitory effect of Sainfoin extracts on L<sub>3</sub> of *H. contortus* and *T. colubriformis* and on adult *T. circumcincta*.

Positive in vivo results have been shown in several studies. Thamsborg et al. (2001) observed in an in vivo study normal growth rates for lambs on Sainfoin, a more than 50% reduced faecal egg output of infected animals and the tendency of lower establishment of incoming larvae and expulsion of adult worm burdens. Recently there have been many studies on the possible effect of Sainfoin with all confirming the significant reduction of FEC after the consumption of Sainfoin (Paolini et al., 2003c; Thamsborg et al., 2003; Athanasiadou et al., 2005b; Hoste et al., 2005a; Lüscher et al., 2005; Paolini et al., 2005b).

There were two short-term trials conducted on the effect of Sainfoin consumption on establishment of incoming larvae in which no evidence for any significant effects could be found (Athanasiadou et al., 2005; Paolini et al., 2005a). These findings are supported by Thamsborg et al. (2003) who showed that the consumption of Sainfoin previous to infection did not influence establishment. The only study on Sainfoin without any significant effects remains the one of Athanasiadou et al. (2005) but they cite that the concentration of CT in the Sainfoin swards might have been insufficient.

The reviewed research results provide evidence that the administration of Sainfoin can reduce faecal egg count and therefore lead to lower pasture contamination. Lüscher et al. (2005) continue to research the potential for the practical integration of tanniferous plants into agricultural practise in their large-scale project and argue that first results in particular on Sainfoin show that this plant has promising potential. In fact not all studies display the same optimism when it comes to the cultivation of Sainfoin. Thamsborg (2001b) states that Sainfoin appears not competitive in grassland swards. Athanasiadou et al. (2005b) also mention poor establishment of Sainfoin in UK.

Frame (2006) writes in about Sainfoin that “monocultures lack competitiveness to weed invasion” when compared to cultivation of mixed Sainfoin/grass stands.

This indicates the need for further research particularly on the cultivation side of feasibility. No recommendation can be given as long as it is not clear how suitable Sainfoin is for different climates and especially for the non chemical methods of organic agriculture.

## **2.5 Quebracho (*Schinopsis ssp.*, family: *Anacardiaceae*)**

Quebracho is the Spanish name for a group of similar species of trees that are originated in tropical South America (Wikipedia, 2006). In the medical context Quebracho is referred to as an extraction from the bark of one the *Schinopsis ssp.* which is rich in condensed tannins (Paolini et al., 2003a).

The first study to be considered in this context in an extensive long-term (10 weeks) feeding study on sheep by Athanasiadou et al. (2000) with several interesting results: the administration of quebracho led to a reduction in FEC and later to lower female fecundity and adult worm burden, however the observed differences were not always significant. During the trial it was also noticed that the animals receiving quebracho had a lower live weight gain and inferior food conversion, although performance of parasitized animals did not decline to the same extent as control animals.

Two short-term experiments were conducted by Paolini et al. (2003ab); in these experiments previously infected goats were drenched with quebracho extract on a daily basis for 8 days. One group was infected with *T. colubriformis* and *T. circumcincta* (2003a) and the other group was infected with *H. contortus*. In all groups the treatment led to a reduction in egg excretion and female fecundity, with no change in the established adult worm population. This study also tested the effect of quebracho on incoming larvae of both species and a reduction was only observed for *T. colubriformis*, the reduction of *T. circumcincta* larvae was insignificant.

The following conclusions can be drawn from the above experimentation: the administration of quebracho extract appears to lead to a reduction of egg excretion and female parasite fecundity but it does not seem to lower the adult worm burden. There are no verified results available on effect of incoming larvae. However studies on quebracho administration remain scarce and although the available results are relatively clear and indicate anthelmintic properties for quebracho, more studies are required to confirm the existing results.

A further critical point that needs to be proved is the availability of quebracho extract, which is certainly also a question of costs. It needs to be estimated whether farms can afford to purchase the delivery of this extract, or whether the application of quebracho will be limited to areas where *Schinopsis ssp.* are naturally grown.

## 2.6 Other plant species with high tanniferous contents

Apart from the above described plants, a range of grasses, shrubs and bushes have been chemically analysed for their CT-content and anthelmintic activity in vitro. Some have shown promising results when tested in vivo however not with the same benefits shown as the previously examined plants (Tab. 1).

The following list shows tanniferous plants that have been the item of scientific consideration, either in in vitro, or in vivo studies, or both. The list is not exhaustive with research in this field an ongoing process.

**Tab. 1. Studies on the anthelmintic effect of different plants/herbs**

Scientific name	Traditional name	Studies
<b>Forage plants:</b>		
<i>Dorycnium pentaphyllum</i>	Socarillo	Molan et al., 2000b Niezen et al., 2002b
<i>Dorycnium rectum</i>	No common name	Molan et al., 2000b, Niezen et al., 2002b Waghorn et al., 2006
<i>Lespedeza cuneata</i>	Chinese Lespedeza	Min and Hart, 2003; Min et al., 2005
<i>Rumex obtusifolius</i>	Dock	Molan et al, 2000b; Thamsborg, 2001a
<b>Shrubs and trees:</b>		
<i>Acacia karoo</i>	Wattle (leaves)	Kahiya et al., 2003
<i>Calluna vulgaris</i>	Heather	Hoste et al., 2005b
<i>Castanea sativa</i>	Chestnut Tree (fruit)	Hoste et al., 2005b
<i>Cornus sanguinea</i>	Common Dogwood	Athanasiadou et al., 2005b
<i>Corylus avellana</i>	Hazel tree	Paolini et al., 2004, Rahmann et al. 2007
<i>Erica ssp.</i>	Erica	Athanasiadou et al., 2005b
<i>Pinus sylvestris</i>	Pine tree (leaves)	Hoste et al., 2005b
<i>Punica granatum</i>	Pomegranate	Athanasiadou et al., 2005b

<i>Quercus ssp.</i>	Oak	Paolini et al., 2004; Hoste et al., 2005b; Athanasiadou et al., 2005b
<i>Robinia pseudacacia</i>	Black Locust	Athanasiadou et al., 2005b
<i>Rubus fruticosus</i>	Blackberry bush	Paolini et al., 2004; Hoste et al., 2005b
<i>Salix ssp.</i>	Willow	Barry et al., 2005; Diaz-Lira et al., 2005
<i>Sarothamnus scoparius</i>	Genista (leaves)	Hoste et al., 2005b
<i>Vitis ssp. extract</i>	Grape Seed extract	Waghorn et al., 2006

Shrubs and trees are of particular interest because of the alimentary spectrum of sheep and goats and their ability to browse a wide range of plants. In this context the nutritional consequences of goats browsing rangeland environments has been extensively studied, but it has taken longer to focus research on the correlation between browsed plant species and parasite infection status. Despite the scarcity of studies there is evidence of a positive effect on parasitism (Hoste et al., 2001; Hoste et al., 2005c). Although sheep are generally categorized as grazers, they still browse trees and shrubs to some extent and as current research results from New Zealand indicate this can have beneficial effects on the resilience to parasitic nematodes (Diaz Lira et al., 2005).

### 3 Phytotherapeutical measures against internal parasites

Phytotherapy is either prophylactical or therapeutical use of plants, their plant components or their preparations, and can be divided into allopathic and traditional phytotherapy (Hördegen, 2005). The allopathic phytotherapeutical approach uses scientific testing to verify the anthelmintic effectiveness of a plant or preparation, and in contrast to that the use of traditional products is based on handed down knowledge (Anonymous, 2005 in: Hördegen, 2005).

The evaluation was focused on possible risks and side-effects of plants, and on scientific verification. Listed are herbs and preparations that have either proved to have an anthelmintic efficacy (scientifically tested) or that are traditionally associated to help against internal parasites (Tab. 2). The criterion for inclusion into the table was the frequency in which plants were mentioned in coherence with anthelmintic activity. It is important to keep in mind when considering alternative options in this area that there still remain a lot of plants not evaluated.

**Tab. 2. Plants and plant preparations used as alternative anthelmintics**

Botanical & Common Name	Scientific Verification & Comment	Prime Source
<i>Allium sativum</i> (Garlic)	traditionally applied In vivo trial showed no effect of garlic administration contradicting statements of the effectiveness	Allen, 1998 Duval, 1994 Cabaret et al., 2002b Meat New Zealand, 1998 Perezgroves, no date University of Aberdeen, no date
<i>Annonum senegalensis</i> (Custard tree)	Traditionally used by Nigerian farmers In vitro test showed promising potential	Alawa et al., 2003
<i>Artemisia abrotanum</i> (Southern Wormwood)	Although traditionally used, activity seems reliable	Hoffmann, 1995 PFAF, 2002
<i>Artemisia absinthium</i> (Common Wormwood)	Reputation of anthelmintic effect in trad. medicine In vitro tests showed barley sign. reduction of <i>Trichstrongylus</i>	Bara et al., 1999 Duval, 1994
<i>Artemisia cina</i> (Eurasian Wormwood)	It is used for the fabrication of Santonin which is used in human medicine	PFAF, 2002 Duval, 1994

<i>Artemisia dracunculus</i> (Tarragon)	Is known to have vermifuge properties in the traditional medicine	Duval, 1994 PFAF, 2002
<i>Artemisia herba-alba</i>	Powder was used in a trial with goats infected with <i>H. contortus</i> and worked successfully	Idris et al., 1982
<i>Artemisia vulgaris</i> (Common Mugwort)	No reliable source could be found that confirms the unobjectionable efficacy	Duval, 1994 PFAF, 2002 www.feenkraut.de
<i>Asarum canadense</i> (Wild Ginger, Snakeroot)	Is known to work as an anthelmintic, traditionally used in Africa, no conducted trial could be found	Duval, 1994 PFAF, 2002
<i>Azadiracta indica</i> (Neem tree)	Results are scientifically verified but vary	Costa et al., 2006 Githiori, 2004 Thomas et al., no date www.neem-foundation.org
<i>Carica papaya</i> (Papaya)	In vitro trials have confirmed anthelmintic activity. Used traditionally in the Philippines and other countries. In vivo trials with papaya seeds in sheep showed 80% reduction in FEC In vivo trial with calves show reduction of 60%	Animal Science at Cornell University, 2001 Hoste et al., 2005d Hounzangbe-Adote et al., 2005a Ronoredjo and Bastiaensen, no date Stepek et al., 2004
<i>Chenopodium ambrosioides</i> (Goosefoot, Wormseed)	Internet database and Cornell University seems to have investigated thoroughly	Animal Science at Cornell University, 2001 PFAF, 2002
<i>Chrysanthemum cinerariifolium</i> (Pyrethrum)	It has been discussed for years but two in vivo trials found no and very low anth. efficacy	Duval, 1994 Hammond et al., 1997 Mbaria et al., 1998
<i>Crucifers</i> (There are several species of the family cuciferae that are used as anthemintics)	All traditional plants with anthelmintical reputation, no in vivo experiments could be found	Duval, 1994 PFAF, 2002
<i>Curcubita pepo</i> (Pumpkin)	Activity scientifically researched, reliable sources	Anonymus, 2003 Duval, 1994 Hoffmann, 1995 PFAF, 2002
<i>Daucus carota</i> (Wild carrot)	Plant can not be recommended because of missing scientific verification	Duval, 1994 PFAF, 2002
<i>Dryopteris ssp.</i> (Fern) ( <i>D. filix-mas</i> quoted the most)	Popular and effective against tapeworms	Cabaret et al., 2002b Duval, 1994 PFAF, 2002
<i>Eucalyptus grandis</i> (Eucalyptus)	In vivo test in goats led to 90% reduced FEC on <i>H. contortus</i> but none on <i>Ostertagia</i>	Animal Science at Cornell University, 2001 Bennet-Jenkins and Bryant, 1996
<i>Ferula conocaula</i> , <i>F. gigantea</i> , <i>F. narthex</i> (Fennel)	Traditionally used, no scientific tests available	Duval, 1994
<i>Fumaria parviflora</i> (Small-flowered/ Fine-leaved Fumitory)	Extract had the same efficiency as common anthelmintic control product Promising alternative	Hördegen et al., 2003 FIBL activity report, 2004
<i>Juglans regia</i> (English Walnut, Black Walnut)	Long history of medical use Traditional anthelm. Known to expel worms	Edward, no date
<i>Khaya senegalensis</i> (Gambian Mahagony)	Traditionally used as a vermifuge In vitro and in vivo tests confirmed anthelmintic potential	Ademola et al., 2004

<i>Mallotus philippensis</i> (Kamala tree)	Scientific trials confirmed in vitro and in vivo anthelmintic activity	Akhtar and Ahmad, 1992 Singh et al., 2004
<i>Melia azedarach</i> (Chinaberry tree) Indian Lilac	Trial with goats was successful and showed virtually no side-effects.	Akhtar and Riffat, 1984 PFAF, 2002
<i>Melinis minutiflora</i> ( <i>Panicum minutiflora</i> ) ( <i>Panicum melinis</i> )	Traditionally used in the Dominican Republic as a de-wormer. Cornell University did in vitro tests on <i>H. contortus</i> that indicated some effect.	Animal Science at Cornell University, 2001
<i>Newbouldia laevis</i> Boundary Tree	Traditionally used by farmers in W.-Africa Confirmation of anthelm. Activity through in vitro testing	Brown, 1992 Hounzangbe-Adote et al., 2005a
<i>Nigella sativa</i> (Black Cumin)	Traditionally used as a anthelmintic, in vivo trial with Monezia confirm effect on tapeworms	Iqbal et al., 2005b PFAF, 2002
<i>Ocimum sanctum</i> (Sacred Basil) <i>Ocimum gratissimum</i> (Basil)	Confirmation of in vitro testes with <i>H. contortus</i>	Anthony et al., 2005 Asha et al., 2001 Pessoa et al., 2002
<i>Spigelia marilandica</i> (Pinkroot, Indian Pink)	Especially effective with tape- and roundworms, treatment should always be followed by a saline laxative	University of Aberdeen, no date PFAF, 2002
<i>Tanacetum vulgare</i> (Tansy)	Traditionally applied, no in vivo results available. In vitro testing showed effectiveness. Other species seem to possess anthelm. properties as well	Duval, 1994 Gadziev and Eminov, 1986 PFAF, 2002
<i>Zanthoxylum zanthoxyloides</i> (Fagara)	Traditionally applied in Western Africa In vitro tests confirmed activity In vivo tests showed that regular feeding is better then a single cure	Hounzangbe-Adote et al., 2005a Hounzangbe-Adote et al., 2005b

At the current stage veterinarians that are willing to work with phytotherapy have to deal with two basic problems. First of all effective plants have to be divided from ineffective ones, and secondly the physiological consequences of herbal administration need to be determined, including possible risks and side-effects. Preparations derived from plants are often thought to be harmless and widely associated to have fewer side effects and are therefore considered easier to apply, however reality shows that plants and plant extracts can be as toxic as allopathics, that they can have side effects, and if applied inappropriately they can cause severe damage and even lead to death (Reichling and Saller, 2001).

A further problem that has been explained and discussed by Häublein (2005) is the legal requirements for the use of homeopathical and phytotherapeutical remedies in the EU. These remain restrictive and discouraging for both veterinarians and farmers, with no change in the foreseeable future. For further information on this matter refer to Häublein (2005). So before advising or suggesting anything to practically working people, these aspects should be well thought of.

#### 4 Copper-oxide wire particles

The basic principal of this treatment is that the availability of macro-minerals and trace elements influences the host-parasite relationship (Suttle and Jones, 1989 in: Chartier et al., 2000). When copper-oxide wire particles (COWP) are administered they remain in the rumen and release free copper into the abomasum which creates an environment that affects *H. contortus* ability to remain established (Burke et al., 2004).

Bang et al. (1990) found that the administration of copper-oxide wire particles led to a good reduction of parasite burden of *H. contortus* but the impact on other species was very average.

Further research proved the efficiency of (COWP) on *H. contortus* in goats but also showed that it does not influence greatly on other species (Chartier et al., 2000).

Other research come to similar conclusions, the treatment seems to successfully reduce FEC and the number of established adults of *H. contortus* but does not work effectively on other species (Watkins, 2003). Burke et al. (2004) evaluated to optimal dosage for administration and found 2g as a single dose to be sufficient to result in reduced FEC and worm burden but not enough to lead to toxicity or predispose lambs to disease which higher concentrations do.

## **5 Homeopathy**

Homeopathy will not be discussed in detail for two reasons. The first is that the last paper that has been written about alternative helminths control a year ago has already dealt with homeopathy sufficiently and the report on homeopathy can be recommended (Häublein, 2005). The second reason is that according to the findings of the last paper homeopathy is considered unsuitable to treat acute helminthosis in most cases. This is due to several circumstances, amongst them the lack of veterinarians that have an additional homeopathic qualification and that the application of homeopathics requires detailed knowledge as incorrect dosing rates can lead to overreaction and worsen the condition.

Despite the above findings it still cannot be generally claimed that it is impossible to de-worm with homeopathic remedies, good results have been obtained in the past in independent reports (Gibbons, 2002).

The mode of action of homeopathy is based on a thorough anamnesis and on the provision of adequate animal husbandry. It requires time and the will of the farmer to think over the whole farming process in order to detect the source of susceptibility.

In conclusion, it can be ascertained that homeopathy has the potential to help the animal to overcome its deteriorating condition caused by parasitic infection but it is currently considered unsuitable as a short-term measure to treat intestinal nematodes on organic farms (Cabaret et al., 2002b; Humann-Ziehank and Ganter, 2005).

## 6 Conclusion

In conclusion future research in this area seems promising, there remain a lot of plants to be tested but it is possible that there are some plants that possess a high anthelmintic effectiveness and cause no side-effects.

In the retrospection of the literature research it has been concerning to discover the incomplete and possible misleading information that is available on the internet and in other sources. The information published by Duval (1994) and the University of Aberdeen is possibly well meant but it can not be considered sufficient to only occasionally and briefly mention possible risks of plants and their preparations. This information may be misinterpreted, with people applying the information in a belief that they are doing their animals a favour by not using common anthelmintics but ending up doing severe damage to their stock.

Some preparations like copper sulphate should actually not even be mentioned as an alternative treatment. So one needs to be very careful before giving any advice to practically working farmers and only introduce plants and preparations with reference to risks and side-effects.

## 7 References

- Abdulrazak SA, Njuguna EG, Karau PK** (2005) The effect of supplementing Rhodes grass (*Chloris gayana*) hay with *Acacia tortilis* leaves and pods mixture on intake, digestibility and growth performance of goats. *Livestock Research for Rural Development* 17
- Ademola IO, Fagbemi BO, Idowu SO** (2004) Evaluation of the anthelmintic activity of *Khaya senegalensis* extract against gastrointestinal nematodes of sheep: in vitro and in vivo studies. *Veterinary Parasitology* 122:151-164
- Aerts RJ, Barry TN, McNabb WC** (1999) Polyphenols and agriculture: beneficial effects of proanthocyanidins in forages. *Agric Ecosystems Environment* 75:1-12
- Alawa CBI, Adamu AM, Gefu JO, Ajanusi OJ, Abdu PA, Chiezey NP, Alawa JN, Bowman DD** (2003) In vitro screening of two Nigerian medicinal plants (*Vernonia amygdalina* and *Annona senegalensis*) for anthelmintic activity. *Veterinary Parasitology* 113:73-81
- Anthony J-P, Fyfe L, Smith H** (2005) Plant active components - a resource for antiparasitic agents? Review in: *Trends in Parasitology*. 21 No10
- Asha MK, Prashanth D, Murali B, Padmaja R, Amit A** (2001) Anthelmintic activity of essential oil of *Ocimum sanctum* and *eugenol*. *Fitoterapia* 72:669-70
- Athanasiadou S, Kyriazakis I, Jackson F, Coop RL** (2000b) Consequences of long-term feeding with tannins on sheep parasited with *T. colubriformis*. *International Journal for Parasitology* 30:1025-1033
- Athanasiadou S, Kyriazakis I, Jackson F, Coop RL** (2001) Direct anthelmintic effects of tannins towards different gastrointestinal nematodes of sheep *in vitro* and *in vivo* studies. *Veterinary Parasitology* 99:205-219
- Athanasiadou S, Tzamaloukas O, Kyriazakis I, Jackson F, Coop RL** (2005) Testing for direct anthelmintic effects of bioactive forages against *Trichostrongylus colubriformis* in grazing sheep. *Veterinary Parasitology* 127:233-243
- Athanasiadou S, Tzamaloukas O, Kyriazakis I, Jackson F, Thamsborg SM, Christensen LP** (2005b) The role of bioactive plants to control sheep nematodes in Northern Europe. In: **Thamsborg SM, Larsen M, Busch M**(eds) (2004) Sustainable, non-chemical control of small ruminant nematode parasites in Europe. Proceedings from an International Workshop held at Danish Centre of Experimental Parasitology Royal Veterinary and Agricultural University
- Bang KS, Familton AS, Sykes AR** (1990) Effect of copper oxide wire particle treatment on establishment of major gastrointestinal nematodes in lambs. *Research in Veterinary Science* 49:132-139
- Bara S, Zaragoza C** et al. (1999) Allelopathic and anthelmintic effect of wormwood (*Artemisia absinthium*). SEMh Congreso 1999: Sociedad Espanola de Malherbologia, Actas Longrono, Spain. In: **Devantier B** (ed) (2004) Evidence for the effectiveness of therapeutic remedies acceptable for the management of internal parasites of sheep and cattle in organic farming systems: A critical review. A paper presented in partial fulfilment of the requirements for a Graduate Diploma of Rural Studies at Massey University, New Zealand

- Barrau E, Fabre N, Fouraste I, Hoste H** (2005) Effect of bioactive compounds from Sainfoin (*Onobrychis viciifolia* Scop.) on the in vitro larval migration of *Haemonchus contortus*: role of tannins and flavonol glycosides. *Parasitology* 131:531-538
- Barry T, Kemp P, Diaz Lira C, McWilliam E, Cameron P, Ravenwood J** (2005) Progress on Wairarapa Farms[online]. New Zealand, zu finden in <[www.hortresearch.co.nz/projects/fodder](http://www.hortresearch.co.nz/projects/fodder)> [cited on 20.4.2006]
- Bennet-Jenkins E, Bryant C** (1996) Novel sources of anthelmintics. *International Journal of Parasitology* 26:937-947
- Brown K** (1992) Medicinal plants, indigenous medicine and conservation of biodiversity in Ghana[online]. United Kingdom, zu finden in <[www.uea.ac.uk/env/cserge/pub/wp/gec/gec\\_1992\\_36.pdf](http://www.uea.ac.uk/env/cserge/pub/wp/gec/gec_1992_36.pdf)> [cited on 27.4.2006]
- Burke JM, Miller JE, Olcott DD, Olcott BM, Terrill TH** (2004) Effect of copper oxide wire particles dosage and feed supplement level on *Haemonchus contortus* infection in lambs *Veterinary Parasitology* 123:235–243
- Chartier C, Etter E, Hoste H, Pors I, Koch C, Dellac B** (2000) Efficacy of Copper Oxide Needles for the Control of Nematode Parasites in Dairy Goats. *Veterinary Research Communications* 24:389-399
- Coop RL, Kyriazakis I** (2001) Influence of host nutrition on the development and consequences of nematode parasitism in ruminants. *Trends in Parasitology* 17:325-330
- Cooper J, Gordon IJ, Pike AW** (2000) Strategies for the avoidance of faeces by grazing sheep. *Applied Animal Behaviour Science* 69:15–33
- Cosgrove GP, Niezen JH** (2000) Intake and selection for white clover by grazing lambs in response to gastrointestinal parasitism. *Applied Animal Behaviour Science* 66:71–85
- Costa CT, Bevilaqua CM, Maciel MV, Camurca-Vasconcelos AL, Morais SM, Monteiro MV, Farias VM, da Silva MV, Souza MM** (2006) Anthelmintic activity of *Azadirachta indica* A. Juss against sheep gastrointestinal nematodes. *Veterinary Parasitology* 137:306-316
- Dawson JM, Buttery PJ, Jenkins D, Wood CD, Gill M** (1999) Effects of dietary quebracho tannin on nutrient utilisation and tissue metabolism in sheep and rats. *Journal of the Science of Food and Agriculture* 79:1423-1430
- Devantier B** (2004) Evidence for the effectiveness of therapeutic remedies acceptable for the management of internal parasites of sheep and cattle in organic farming systems: A critical review. A paper presented in partial fulfilment of the requirements for a Graduate Diploma of Rural Studies at Massey University, New Zealand
- Diaz Lira C, Barry T, Pomroy B, McWilliam E** (2005) Controlling parasites in weaned lambs on browse blocks[online]. New Zealand, zu finden in <[www.hortresearch.co.nz/index/page/549](http://www.hortresearch.co.nz/index/page/549)> [cited on 20.4.2006]
- Duval J** (1994) The control of internal parasites in ruminants[online]. Canada, zu finden in <[www.eap.mcgill.ca/AgroBio/ab370-04e.htm#BOTANICAL%20DEWORMERS](http://www.eap.mcgill.ca/AgroBio/ab370-04e.htm#BOTANICAL%20DEWORMERS)> [cited on 19.3.2006]
- Edward F** (no date) Black Walnut Hull[online]. USA, zu finden in <[www.parabuster.com/learning/black-walnut-hull.php](http://www.parabuster.com/learning/black-walnut-hull.php)>
- Frame J, Charlton JFL, Laidlaw AS** (1998) Temperate Forage Legumes. Wallingford : CAB International, 327 p
- Frame J** (2006) *Hedysarum coronarium* L.. FAO-database[online]. Zu finden in <<http://www.fao.org/ag/agp/agpc/doc/gbase/data/>>
- Gadzhiev YG, Eminov RS** (1986) Action of medical plants on gastro-intestinal nematodes of sheep. Byulleten Vsesoyuznogo Instituta Gel'minthologii im K.I. Skryabina. In: **Devantier B**(ed) (2004) Evidence for the effectiveness of therapeutic remedies acceptable for the management of internal parasites of sheep and cattle in organic farming systems: A critical review. A paper presented in partial fulfilment of the requirements for a Graduate Diploma of Rural Studies at Massey University, New Zealand
- Gibbons J** (2002) Livestock Research. Alternative methods of Internal Parasite Control in Sheep[online]. United Kingdom, zu finden in <[www.abdn.ac.uk/organic/organic\\_34.php](http://www.abdn.ac.uk/organic/organic_34.php)> [cited on 30.4.2006]
- Githiori JB** (2004) Evaluation of Anthelmintic Properties of Ethnoveterinary Plant Preparations Used as Livestock Dewormers by Pastoralists and Small Holder Farmers in Kenya. Doctoral thesis. Swedish University of Agricultural Sciences, Uppsala
- Hammond JA, Fielding D, Bishop SC** (1997) Prospects for plant anthelmintics in tropical veterinary medicine. *Veterinary Research Communications* 21:213-228
- Häublein K** (2005) Alternative Endo-Parasitenkontrolle bei Schafen. Diplomarbeit. Fachbereich Ökologische Agrarwissenschaften, Universität Kassel
- Heckendorn F** (2005) Kondensierte Tannine - Eine Möglichkeit zur Kontrolle von Magen-Darm-Würmern? *Forum* 1/2, pp 11-13
- Hördegen P** (2005) Epidemiology of internal parasites on Swiss organic dairy farms and phytotherapy as a possible worm control strategy. Dissertation. Swiss Federal Institute of Technology Zürich

- Hördegen P, Hertzberg H, Heilmann J, Langhans W, Maurer V** (2003) The anthelmintic efficacy of five plant products against gastrointestinal trichostrongylids in artificially infected lambs. *Veterinary Parasitology* 117:51–60
- Hoste H, Athanasiadou S, Paolini V, Jackson F, Valderrábano J** (2004) Nutritional aspects of bioactive forages for nematode control in organic sheep and goats. In: **Hovi M, Sundrum A, Padel S** (eds) (2004) Organic livestock farming: potential and limitations of husbandry practice to secure animal health and welfare and food quality. Proceedings of the 2<sup>nd</sup> SAFO Workshop, 25-27 March 2004, Witzenhausen, Germany.
- Hoste H, Gaillard L, Le Frileux Y** (2005a) Consequences of the regular distribution of sainfoin hay on gastrointestinal parasitism with nematodes and milk production in dairy goats. *Small Ruminant Research* 59:265-271
- Hoste H, Lévesque H, Dorchies Ph** (2001) Comparison of nematode infections of the gastrointestinal tract in Angora and dairy goats in a rangeland environment: relations with the feeding behaviour. *Veterinary Parasitology* 101:127-135
- Hoste H, Paolini V, Valderrabano J, Uriarte J, Barrau E, Fouraste I** (2005b) Use of bioactive plants to control infections of the gastrointestinal tract with nematodes in goats in the Southern part of Europe. In: **Thamsborg SM, Larsen M, Busch M** (eds) (2004) Sustainable, non-chemical control of small ruminant nematode parasites in Europe. Proceedings from an International Workshop held at Danish Centre of Experimental Parasitology Royal Veterinary and Agricultural University
- Hoste H, Torres-Acosta JF, Paolini V, Aguilar-Caballero A, Etter E, Lefrileux Y, Chartier C, Broqua C** (2005c) Interactions between nutrition and gastrointestinal infections with parasitic nematodes in goats. *Small Ruminant Research* 60:141-151
- Hoste H, Hounzangbe-Adote S, Fouraste I, Moutairou K** (2005d) In vitro effects of four tropical plants on the activity and development of the parasitic nematode, *Trichostrongylus colubriformis*. *Journal of Helminthology* 79:29-33
- Hounzangbe-Adote MS, Paolini V, Fouraste I, Moutairou K, Hoste H** (2005a) In vitro effects of four tropical plants on three life-cycle stages of the parasitic nematode, *Haemonchus contortus*. *Research in Veterinary Science* 78:155-160
- Hounzangbe-Adote MS, Zinsou FE, Hounpke V, Moutairou K, Hoste H** (2005b) In vivo effects of Fagara leaves on sheep infected with gastrointestinal nematodes. *Tropical Animal Health and Production* 37:205-214
- Iqbal Z, Jabbar A, Akhtar MS, Muhammad G, Lateef M** (2005) Possible Role of Ethnoveterinary Medicine in Poverty Reduction in Pakistan: Use of Botanical Anthelmintics as an Example. *Journal of Agriculture and Social Science* 2:187-195
- Jackson F** (2005) Other control measures. In: **Thamsborg SM, Larsen M, Busch M** (eds) (2004). Sustainable, non-chemical control of small ruminant nematode parasites in Europe. Proceedings from an International Workshop held at Danish Centre of Experimental Parasitology Royal Veterinary and Agricultural University
- Kahiya C, Mukaratirwa S, Thamsborg SM** (2003) Effects of *Acacia nilotica* and *A. karoo* diets on *Haemonchus contortus* infection in goats. *Veterinary Parasitology* 115:265-274
- Kassai T** (1999) *Veterinary Helminthology*. Oxford : Butterworth-Heinemann
- Lolicato S** (1998) Lotus. Agriculture Notes[online]. Australia, zu finden in [www.dpi.vic.gov.au/dpi/nreninf.nsf/9e58661e880ba9e44a256c640023eb2e/37a0371eff4a8c2dca256f0f0020f3af/\\$FILE/AG0718.pdf](http://www.dpi.vic.gov.au/dpi/nreninf.nsf/9e58661e880ba9e44a256c640023eb2e/37a0371eff4a8c2dca256f0f0020f3af/$FILE/AG0718.pdf) [cited on 19. April 2007]
- Lüscher A, Häring DA, Heckendorn F, Scharenberg A, Dohme F, Maurer V, Hertzberg H** (2005) Use of tanniferous plants against gastro-intestinal nematodes in ruminants. In: Researching Sustainable Systems - International Scientific Conference on Organic Agriculture, Adelaide, Australia, September 21-23
- Marley CL, Cook R, Barrett J, Keatinge R, Lampkin NH, McBride SD** (2003) The effect of dietary forage on the development and survival of helminth parasites in ovine faeces in: *Veterinary Parasitology* 118:93-107
- Marley CL, Cook R, Keatinge R, Barrett J, Lampkin NH** (2003a) The effect of birdsfoot trefoil (*Lotus corniculatus*) and chicory (*Cichorium intybus*) on parasite intensities and performance of lambs naturally infected with helminth parasites. *Veterinary Parasitology* 112:147–155
- Marley CL, Fraser MD, Fychan R, Theobald VJ, Jones R** (2005) Effect of forage legumes and anthelmintic treatment on the performance, nutritional status and nematode parasites of grazing lambs. *Veterinary Parasitology* 131:267-282
- Mbaria JM, Maitho TE, Mitema ES, Muchiri D J** (1998) Comparative efficacy of pyrethrum marc with albendazole against sheep gastrointestinal nematodes. *Tropical Animal Health and Production* 30:17-22
- Min BR, Hart SP, Miller D, Tomita GM, Loetz E, Sahlu T** (2005) The effect of grazing forage containing condensed tannins on gastro-intestinal parasite infection and milk composition in Angora does. *Veterinary Parasitology* 130:105-113
- Min BR, Hart SP** (2003) Tannins for suppression of internal parasites. *Journal of Animal Science* 81:102-109

- Molan AL, Alexander RA, Brookes IM, MacNabb WC** (2000a) Effect of an extract from Sulla (*Hedysarum coronarium*) containing tannins on the migration of three sheep gastrointestinal nematodes in vitro. *Proceedings of New Zealand Society of Animal Production* 60:21-25
- Molan AL, Waghorn GC, Min BR, McNabb WC** (2000b) The effect of condensed tannins from seven herbage on *Trichostrongylus colubriformis* larval migration in vitro. *Folia Parasitologica* 47:39-44
- Nfi AN, Mbanya JN, Ndi C, Kameni A, Vabi M, Pingpoh D, Yonkeu S, Moussa C** (2001) Ethnoveterinary Medicine in the Northern Provinces of Cameroon. *Veterinary Research Communications* 25:71-76
- Niezen JH, Charleston WAG, Robertson HA, Shelton D, Waghorn GC, Green R** (2002a) The effect of feeding sulla (*Hedysarum coronarium*) or lucerne (*Medicago sativa*) on lamb parasite burdens and development of immunity to gastrointestinal nematodes. *Veterinary Parasitology* 105:229-245
- Niezen JH, Robertson HA, Sidey A, Wilson SR** (2002c) The effect of pasture species on parasitism and performance of lambs grazing one of three grass-white clover pasture swards. *Veterinary Parasitology* 105:303-315
- Niezen JH, Robertson HA, Waghorn GC, Charleston WAG** (1998) Production, faecal egg counts and worm burdens of ewe lambs which grazed six contrasting forages. *Veterinary Parasitology* 80:15-27
- Niezen JH, Waghorn GC, Graham T, Carter JL, Leathwick DM** (2002b) The effect of diet fed to lambs on subsequent development of *Trichostrongylus colubriformis* larvae in vitro and on pasture. *Veterinary Parasitology* 105:269-283
- Paolini V, Fouraste I, Hoste H** (2004) In vitro effects of three woody plant and sainfoin extracts on two parasitic stages of three parasitic nematode species. *Parasitology* 129:69-77
- Paolini V, Bergeaud JP, Grisez C, Prevot F, Dorchies Ph, Hoste H** (2003b) Effects of tannins on goats experimentally infected with *Haemonchus contortus*. *Veterinary Parasitology* 113:253-261
- Paolini V, De La Farge F, Prevot F, Dorchies Ph, Hoste H** (2005b) Effects of the repeated distribution of sainfoin hay on the resistance and the resilience of goats naturally infected with gastrointestinal nematodes. *Veterinary Parasitology* 127:277-283
- Paolini V, Dorchies Ph, Hoste H** (2003c) Effects of sainfoin hay on gastrointestinal nematode infections in goats. *Veterinary Record* 152:600-601
- Paolini V, Frayssines A, De La Farge F, Dorchies Ph, Hoste H** (2003a) Efficacy of tannins on established populations and on incoming larvae of *Trichostrongylus colubriformis* and *Teladorsagia circumcincta* in goats. *Veterinary Research* 34:331-339
- Paolini V, Prevot F, Dorchies Ph, Hoste H** (2005a) Lack of effects of quebracho and sainfoin hay on incoming third-stage larvae of *Haemonchus contortus* in goats. Short communication. *The Veterinary Journal* 170:260-263
- Pessoa LM, Morais SM, Bevilaqua CM, Luciano JH** (2002) Anthelmintic activity of essential oil of *Ocimum gratissimum* Linn. and eugenol against *Haemonchus contortus*. *Veterinary Parasitology* 109:59-63
- Pomroy WE, Adlington BA** (2006) Efficacy of short-term feeding of sulla (*Hedysarum coronarium*) to young goats against a mixed burden of gastrointestinal nematodes. Short communication. *Veterinary Parasitology* 136:363-366
- Rahmann G** (2004) Gehölzfutter - eine neue Quelle für die ökologische Tierernährung. *Landbauforsch Völkenrode SH* 272:29-42
- Rahmann G, Koopmann R, Gutperlet J** (2007) The effect of feeding hazelnut leaves on egg output of gastrointestinal-parasites of goats. *Proceedings of the 9. Wissenschaftstagung Ökologischer Landbau* 21.-23. März 2007 in Stuttgart
- Rahmann G, Koopmann R, Hertzberg H** (2002) Gesundheit erhalten statt Krankheit kurieren. FORSCHUNGSReport, Verbraucherschutz, Ernährung, Landwirtschaft. *Forschungs Report Nr. 1* pp 4-7
- Reichling J, Saller R** (2001) Pflanzliche Arzneimittel in der Veterinärphytotherapie. *Schweiz. Arch. Tierheilk.* 143:395-403
- Ronoredjo EP, Bastiaensen PXM** (No Date) The Use of Indigenous Papaya (*Carica papaya*) as an Anthelmintic for the Treatment of Gastro-intestinal Nematodes in Naturally Infected Calves in Suriname[online]. Suriname, zu finden in <[www.bastiaensen.be/11-Ronoredjo.pdf](http://www.bastiaensen.be/11-Ronoredjo.pdf)> [cited on 25.4.2006]
- Shaik SA, Terrill TH, Miller JE, Kouakou B, Kannan G, Kallu RK, Mosjidis JA** (2004) Effects of feeding *Sericea lespedeza* hay to goats infected with *Haemonchus contortus*. *South African Journal of Animal Science* 34:248-250
- Stepek G, Behnke JM, Buttle DJ, Duce IR** (2004) Natural plant cysteine proteinases as anthelmintics? *Trends in Parasitology* 20 No.7
- Sykes AR, Coop RL** (2001) Interactions between nutrition and gastrointestinal parasitism in sheep. *New Zealand Veterinary Journal* 49:222-226
- Thamsborg SM** (2001b) Options for parasite control using tanniferous forages in livestock in Northern temperate areas. Lecture given on: The 18<sup>th</sup> International Conference of the World Association for the Advancement of Veterinary Parasitology: Promoting Advancement, Preserving Tradition, Stresa, Italy, 26.-30. August 2001[online]. Italien, zu finden in <<http://orprints.org/4519>> [cited on 19.4.2006]

- Thamsborg SM** (2001a) Parasite control on organic sheep farms - options and limitations. In: **Hovi M, Vaarst M** (eds) Positive health - Preventive measures and alternative strategies. Proceedings of the Fifth NAHWOA Workshop. Rødning, Denmark, November 11-13
- Thamsborg SM, Larsen M, Busch M** (2004) Sustainable, non-chemical control of small ruminant nematode parasites in Europe. Proceedings from an International Workshop held at Danish Centre of Experimental Parasitology Royal Veterinary and Agricultural University
- Thamsborg SM, Mejer H, Bandier M, Larsen M** (2003) Influence of different forages on gastrointestinal nematode infections in grazing lambs. Lecture given on: The 19th International Conference for the Advancement of Veterinary Parasitology, New Orleans, Louisiana, USA, 10.-14. August 2003[online]. USA, zu finden in <<http://orgprints.org/4520>> [cited on 19.4.2006]
- Thamsborg SM, Mejer H, Roepstorff A** (2001) Sainfoin reduces the establishment of nematode infections in grazing lambs. Lecture given on: The 18th International Conference of the World Association for the Advancement of Veterinary Parasitology: Promoting Advancement, Preserving Tradition, Stresa, Italy, 26.-30. August 2001[online]. Italy, zu finden in <<http://orgprints.org/4517>> [cited on 19.4.2006]
- Thamsborg SM, Roepstorff A** (2003) Parasite problems in organic livestock and options for control. *Journal of Parasitology* 89:277-284
- Thamsborg SM, Roepstorff A, Larsen M** (1999) Integrated and biological control of parasites in organic and conventional production systems. *Veterinary Parasitology* 84:169-186
- Thomas GW, Cooper B, Lauckner B** (no date) Preliminary investigation into the use of neem as an anthelmintic for the control of gastro-intestinal parasites of small ruminants[online]. Antigua, zu finden in <[www.agriculture.gov.bb/files/Neem.pdf](http://www.agriculture.gov.bb/files/Neem.pdf)> [cited on 24.4.2006]
- Tzamaloukas O, Athanasiadou S, Kyriazakis I, Jackson F, Coop RL** (2005) The consequences of short-term grazing of bioactive forage on established adult and incoming larvae populations of *Teledorsagia circumcincta* in lambs. *International Journal for Parasitology* 35:329-335
- Vlassoff A, McKenna PB** (1994) Nematode parasites of economic importance in sheep in New Zealand. *New Zealand Journal of Zoology* 21:1-8
- Waller P.J, Thamsborg SM** (2004) Nematode control in 'green' ruminant production systems. *Trends in Parasitology* 20 No.10
- WatkinsAD** (2003) Effectiveness of copper-wire particles on the control of *Haemonchus contortus* in sheep. A Thesis. Graduate Faculty of the Louisiana State University and Agricultural and Mechanical College