

DO WILDFLOWER STRIPS ENHANCE PEST CONTROL IN ORGANIC

CABBAGE?

J. Schied¹, A. Leimgruber^{2,3}, M. Willareth^{2,3}, P. Nagel³, L. Pfiffner², H. Luka², E. Wyss², C. Schlatter², & M. Traugott¹

¹ Mountain Agriculture Research Unit, Institute of Ecology, University of Innsbruck, Technikerstrasse 25, 6020 Innsbruck, Austria, Michael.Traugott@uibk.ac.at; ² Research Institute of Organic Agriculture, Ackerstraße, 5070 Frick, Switzerland; ³ Institute of Biogeography, Department of Environmental Sciences, University of Basel, St. Johannis-Vorstadt 10, 4056 Basel, Switzerland

Within this project we assess whether wildflower strips and companion plants increase the control of cabbage pests *Plutella xylostella* L. (Lepidoptera: Plutellidae), *Mamestra brassicae* L. (Lepidoptera: Noctuidae) and *Pieris rapae* L. (Lepidoptera: Pieridae) by (1) naturally occurring parasitoids and predators and (2) mass-released *Trichogramma brassicae* (Bezdenko) (Hymenoptera: Trichogrammatidae) parasitoids. Two organic cabbage fields were used for this study: adjacent to each field a wildflower strip was sown and companion plants (*Centaurea cyanus* L. (Asteraceae)) intermixed within the crop. Within each field ~15,000 *M. brassicae* eggs were placed out to determine the parasitism rates by mass-released *T. brassicae* and to assess the levels of egg predation. Over 1,000 lepidopteran larvae were collected and screened for hymenopteran and tachinid parasitoid DNA using a multiplex PCR assay. Invertebrate generalist predators (n=1,063) were collected for DNA-based gut content analysis. The wildflower strip had a significant positive effect on *M. brassicae* egg parasitism rates as rates increased 5-fold in the vicinity to the strip. Moreover, companion plants enhanced invertebrate predation on *M. brassicae* eggs. Both, the release of *T. brassicae* and the use of companion plants, however, did not significantly increase egg parasitism rates. The infestation of plants by caterpillars increased with distance to the wildflower strip and there was a trend of decreasing larval parasitism rates with distance to the strip. Currently the invertebrate predators are being molecularly analysed to assess predation on unparasitized and parasitized lepidopteran pests.