

# Literaturliste

## Kastration bei Schweinen

# Ebermast – eine Herausforderung für den Ökolandbau

Von Mirjam Holinger und Barbara Früh

Artikel erschienen in:

ÖKOLOGIE & LANDBAU 166,  
2/2013, Seite 42–44

## Literatur

- Anonym (2010): **Europäische Erklärung über Alternativen zur chirurgischen Kastration bei Schweinen.** Abrufbar unter [http://ec.europa.eu/food/animal/welfare/farm/docs/castration\\_pigs\\_declaration\\_de.pdf](http://ec.europa.eu/food/animal/welfare/farm/docs/castration_pigs_declaration_de.pdf)
- Baes, C., H. Luther, S. Ampuero, P. Spring, A. Hofer (2011): **The effect of androstenone, skatole and indole on semen quality.** In proceedings of: Boars heading for 2018, 16. Wageningen University and Research Centre, Amsterdam
- Blanch, M., N. Panella-Riera, P. Chevillon, M. F. Furnols, M. Gil, J. M. Gil, Z. Kallas, M. A. Oliver (2012): **Impact of consumer's sensitivity to androstenone on acceptability of meat from entire male pigs in three European countries: France, Spain and United Kingdom.** Meat Science 90, S. 572–578
- Boyle, L. A., L. Björklund (2007): **Effects of fattening boars in mixed or single sex groups and split marketing on pig welfare.** Animal Welfare 16, S. 259–262
- Cronin, G. M., F. R. Dunshea, K. L. Butler, I. McCauley, J. L. Barnett, P. H. Hemsworth (2003): **The effects of immuno- and surgical-castration on the behaviour and consequently growth of group-housed, male finisher pigs.** Applied Animal Behaviour Science 81, S. 111–126
- Doran, E., F. W. Whittington, J. D. Wood, J. D. McGivan (2002): **Cytochrome P4501IE1 (CYP2E1) is induced by skatole and this induction is blocked by androstenone in isolated pig hepatocytes.** Chemico-Biological Interactions 140, S. 81–92
- Fredriksen, B., B. M. Lium, C. H. Marka, B. T. Heier, E. Dahl, J. U. Choinski, O. Nafstad (2006): **Entire male pigs in a farrow-to-finish system. Effects on androstenone and skatole.** Livestock Science 102, S. 146–154

- Fredriksen, B., B. M. Lium, C. H. Marka, B. Mosveen, O. Nafstad (2008): **Entire male pigs in farrow-to-finish pens. Effects on animal welfare.** Applied Animal Behaviour Science 110, S. 258–268
- Frieden, L., C. Looft, D. Mörlein, L. Meier-Dinkel, E. Tholen (2011): **Breeding for reduced boar taint in Pietrain sired crossbred males in Germany.** In proceedings of: Boars heading for 2018, 13. Wageningen University and Research Centre, Amsterdam
- Hansen, L. L., H. Mejer, S. M. Thamsborg, D. V. Byrne, A. Roepstorff, A. H. Karlsson, J. Hansen-Møller, M. T. Jensen, M. Tuomola (2006): **Influence of chicory roots (*Cichorium intybus* L.) on boar taint in entire male and female pigs.** Animal Science 82, S. 359–368

- Jensen, M. T., L. L. Hansen (2006): Feeding with chicory roots reduces the amount of odorous compounds in colon and rectal contents of pigs. Animal Science 82, S. 369–376
- Malmfors, B., K. Lundström (1983): Consumer reactions to boar meat. A review. Livestock Production Science 10, S. 187–196
- Marty, D. (2011): 30'000 Eber pro Woche. KAGMagazin 5, S. 4–6
- Meyer, E. (2011): Was leisten die Eber? Landesamt für Umwelt, Landwirtschaft und Geologie, Köllitsch. Abrufbar unter [www.landwirtschaft.sachsen.de/landwirtschaft/download/MeyerEberleistung\\_Fachinfo.pdf](http://www.landwirtschaft.sachsen.de/landwirtschaft/download/MeyerEberleistung_Fachinfo.pdf)
- Preinerstorfer, A., A. Leithold, G. Huber, B. Krimberger, I. Mösenbacher-Molterer (2010): Erfahrungen zur Ebermast. In: Nutztierschutztagung Raumberg-Gumpenstein 2010. Hrsg. vom Lehr- und Forschungszentrum für Landwirtschaft (LFZ) Raumberg-Gumpenstein. LFZ, Raumberg-Gumpenstein, S. 47–54
- Rydhmer, L., G. Zamaratskaia, H. K. Andersson, B. Algers, R. Guillemet, K. Lundström (2006): Aggressive and sexual behaviour of growing and finishing pigs reared in groups, without castration. Acta Agriculturae Scandinavica, Section A, Animal Science 56, S. 109–119
- Squires, E. J. (2006): Possibilities for selection against boar taint. Acta Veterinaria Scandinavica 48, S. 1–4
- Stoll, P. (2002): Jungebermast in einigen europäischen Ländern. Eidgenössische Forschungsanstalt für Nutztiere, Posieux. Abrufbar unter [www.agroscope.admin.ch/publikationen/einzelpublikation/index.html?lang=de&aid=16708&pid=17438](http://www.agroscope.admin.ch/publikationen/einzelpublikation/index.html?lang=de&aid=16708&pid=17438)

- Vhile, S. G., N. P. Kjos, H. Sørum, M. Øverland (2012): Feeding Jerusalem artichoke reduced skatole level and changed intestinal microbiota in the gut of entire male pigs. Animal 6, S. 807–814
- Xue, J., G. D. Dial, J. E. Pettigrew (1997): Performance, carcass, and meat quality advantages of boars over barrows: A literature review. Swine Health and Production 5, S. 21–28
- Zamaratskaia, G., J. Babol, H. K. Andersson, K. Andersson, K. Lundström (2005): Effect of live weight and dietary supplement of raw potato starch on the levels of skatole, androstenone, testosterone and oestrone sulphate in entire male pigs. Livestock Production Science 93, S. 235–243