

Acorns for fattening free-range pigs

Problem

Montanera is a traditional feeding system where pigs forage for acorns. Iberian pigs foraging for acorns during their finishing phase has economically contributed to conserving the “dehesa”, a high nature value (HNV) farmland based on agroforestry, currently consisting of more than four million hectares in the southwest of the Iberian Peninsula (photo 2).

Solution

This natural resource is used to fatten pigs without any kind of compound or supplementary feed. During the last 2-3 months of fattening, pigs can gain more than 40 kg of body weight from grass and foraged acorns.

Benefits

In the montanera system, the mean average daily gain for fattening pigs is ≥ 0.75 kg/day. Their body fat has a high concentration of oleic acid (around 55%) and very low concentrations of linoleic and palmitic acids; which is very important for the quality of pork and the cured products.

Applicability box

Theme

Pigs, feeding and ration planning

Context

South West of Iberian Peninsula; partly adaptable to other Mediterranean areas and forests with *Quercus* species (the best species is *Q. ilex rotundifolia*).

Application time

Autumn and winter.

Required time

None if there are adult trees; approximately 15 years to have the first mast of acorns if it is necessary to establish trees and the surface is certified as organic.

Period of impact

1.5 months to influence meat quality and fatty acid profile

Equipment

None for free grazing; only a stick to knock down acorns if there is a swineherd with the pigs.

Best in

Fattening pigs (especially in fatty breeds)

Practical recommendation

- The fattening performance is very much influenced by the age of pigs and their compensatory growth; hence, pigs should be as old as possible (≥ 1 year) and adapted to grazing.
- Grass is necessary as a source of protein to compensate for the low protein levels in acorns.
- The food conversion rate is 10.5 kg of whole acorns of *Q. i. rotundifolia* to gain 1 kg, besides the contribution of grass; to establish the stocking rate, consider that an adult evergreen oak produces ≈ 11 kg of acorns/year).
- Iberian pigs peel acorns to avoid the high content of tannins in the shell. However, during peeling, approximately 20% of the kernel can be wasted.




Photo 1: Pig foraging acorns (Vicente Rodríguez-Estévez, University of Córdoba)



Photo 2: Group of Iberian pig fatteners grazing in a dehesa estate (Vicente Rodríguez-Estévez, University of Córdoba)

Further information

Video

- The video [“Cerdos Comiendo Bellotas en la Dehesa”](#) shows a swineherd knocking down acorns.
- The video [“Cerdos ibéricos comiendo bellotas en una dehesa de Extremadura”](#) shows a herd of Iberian pigs foraging acorns.

Further reading

López-Bote, Clemente J. (1998). Sustained utilization of the Iberian pig breed. In: *Meat Science*, Vol. 49, No. Suppl. I, 2018, pp. 17-27, [https://doi.org/10.1016/S0309-1740\(98\)90036-5](https://doi.org/10.1016/S0309-1740(98)90036-5)

Rodríguez-Estévez, Vicente et al. (2007). Producción de bellota en la dehesa: factores influyentes. In: *Archivos de Zootecnia*, Vol.56(R), 2007, pp. 25-43.

Rodríguez-Estévez, Vicente et al. (2008). Dimensiones y características nutritivas de las bellotas de los Quercus de la dehesa. In: *Archivos de Zootecnia*, Vol. 57(R), 2008, pp. 1-12.

Rodríguez-Estévez, Vicente et al. (2009). Intrinsic factors of acorns that influence the efficiency of their consumption by Iberian pigs. In: *Livestock Science*, Vol.122, 2009, pp. 281–285, <https://doi.org/10.1016/j.livsci.2008.09.011>

Rodríguez-Estévez, Vicente et al. (2010). Feed conversion rate and estimated energy balance of free grazing Iberian pigs. In: *Livestock Science*, Vol.132, 2010, pp. 152–156, <https://doi.org/10.1016/j.livsci.2010.05.019>

Rodríguez-Estévez, Vicente et al. (2012). Consumption of Acorns by Finishing Iberian Pigs and Their Function in the Conservation of the Dehesa Agroecosystem. In: *Agroforestry for Biodiversity and Ecosystem Services - Science and Practice*, Martin Leckson Kaonga, IntechOpen, DOI: 10.5772/34877. Available from: <https://www.intechopen.com/books/agroforestry-for-biodiversity-and-ecosystem-services-science-and-practice/consumption-of-acorns-by-finishing-iberian-pigs-and-their-function-in-the-conservation-of-the-dehesa>

Weblinks

- Further documents can be found on the [Organic Farm Knowledge](#) website.

About this practice abstract and OK-Net EcoFeed

Publishers

Asociación Valor Ecológico – Ecovalia, Avenida Diego Martínez Barrio 10, primera planta, modulo 12, 41013 Sevilla, Spain, info@ecovalia.org, www.ecovalia.org

Universidad de Córdoba, Campus Universitario de Rabanales, Departamento de Producción Animal, Facultad de Veterinaria, ES-14071 Córdoba, Spain, www.uco.es

Research Institute of Organic Agriculture (FiBL)
Ackerstrasse 113, Postfach 219, CH-5070 Frick
Phone +41 62 865 72 72, info.suisse@fibl.org, www.fibl.org

IFOAM EU, Rue du Commerce 124, BE-1000 Brussels
Phone +32 2 280 12 23, info@ifoam-eu.org, www.ifoam-eu.org

Authors: Vicente Rodríguez-Estévez, Cipriano Díaz-Gaona, Santos Sanz-Fernández, Carolina Reyes-Palomo, Manuel Sánchez-Rodríguez. All Cátedra de Ganadería Ecológica Ecovalia, (Universidad de Córdoba)

Review: Barbara Früh, FiBL, Switzerland, and Lindsay Whitstance, Organic Research Centre, UK



Contact: vrestevez@uco.es

Permalink: [Organic-farmknowledge.org/tool/37476](https://organic-farmknowledge.org/tool/37476)

OK-Net EcoFeed: This practice abstract was elaborated in the Organic Knowledge Network on Monogastric Animal Feed project. The project is running from January 2018 to December 2020. The overall aim of OK-Net EcoFeed is to help farmers, breeders and the organic feed processing industry in achieving the goal of 100% use of organic and regional feed for monogastrics.

Project website: ok-net-ecofeed.eu

Project partners: IFOAM EU Group (project coordinator), BE; Aarhus University (ICROFS), DK; Organic Research Centre (ORC), UK; Institut Technique de l'Agriculture Biologique (ITAB), FR; Research Institute of Organic Agriculture (FiBL), CH; Bioland, DE; Associazione Italiana per l'Agricoltura Biologica (AIAB), IT; Donau Soja DS, AT; Swedish University of Agricultural Sciences, SE; ECOVALIA, ES; Soil Association, UK.

© 2020

