

# Tropentag 2023

International Research on Food Security, Natural  
Resource Management and Rural Development

## Competing pathways for equitable food systems transformation: trade-offs and synergies

Book of abstracts

**Editor:** Eric Tielkes

**Reviewers/scientific committee:** Ayobami Adetoyinbo, Folkard Asch,  
Christian Bateki Adjogo, Bonnie Blaimer, Michael Brüntrup,  
Robert Cárcamo Mallen, Tsu-Wei Chen, Michelle Chevelev-Bonatti,  
Claudia Coral, Teresa Da-Silva-Rosa, Emmanuel Donkor, Christoph Gornot,  
Stef De Haan, Caroline Hambloch, Harry Hoffmann, Gudrun Keding,  
Marcos Lana, Katharina Lohr, Dagmar Mithöfer, Janvier Ntwali, Regina Rößler,  
Constanze Rybak, Lilli Scheiterle, Barbara Schröter, Johannes Schuller,  
Verena Seufert, Stefan Sieber, Jonathan Steinke, Silke Stöber, Götz Uckert,  
Martin Wiehle, Stefan Winter

**Editorial assistance:** Janna Pfister

# Impressum

Bibliografische Information der Deutschen Nationalbibliothek

Die Deutsche Nationalbibliothek verzeichnet diese Publikation in der Deutschen Nationalbibliografie; detaillierte bibliografische Daten sind im Internet über <http://dnb.ddb.de> abrufbar.

1. Aufl. - Göttingen: Cuvillier, 2023

Tropentag 2023: Competing pathways for equitable food systems transformation: trade-offs and synergies Tielkes, E. (ed.) - Witzenhausen, DITSL

© DITSL

Steinstrasse 19, 37213 Witzenhausen

Telefon: 05542-6070

<https://www.ditsl.org>

Alle Rechte vorbehalten. Ohne ausdrückliche Genehmigung des Verlages ist es nicht gestattet, das Buch oder Teile daraus auf fotomechanischem Weg (Fotokopie, Mikrokopie) zu vervielfältigen.

The authors of the articles are solely responsible for the content of their contribution.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means without prior permission of the copyright owners.

Online-Version: <http://www.tropentag.de/>

© CUVILLIER VERLAG, Göttingen 2023

Nonnenstieg 8, 37075 Göttingen

Telefon: 0551-54724-0

Telefax: 0551-54724-21

[www.cuvillier.de](http://www.cuvillier.de)

1. Auflage, 2023

Gedruckt auf umweltfreundlichem, säurefreiem Papier aus nachhaltiger Forstwirtschaft.

ISBN 978-3-7369-7880-5

eISBN 978-3-7369-6880-6

## How do composting materials and duration impact compost evolution and quality in the tropics?

FELIX MATHERI<sup>1</sup>, ANNEKELLY KAMBURA<sup>2</sup>, MAINA MWANGI<sup>3</sup>, EDWARD KARANJA<sup>1</sup>,  
NOAH ADAMTEY<sup>4</sup>, NANCY MUNYOKI<sup>1</sup>, EDWIN MWANGI<sup>1</sup>, DAVID BAUTZE<sup>4</sup>,  
STEVEN RUNO<sup>3</sup>

<sup>1</sup>*International Centre of Insect Physiology and Ecology (icipe), Kenya*

<sup>2</sup>*Taita Taveta University, Kenya*

<sup>3</sup>*Kenyatta University, Agricultural Science and Technology, Kenya*

<sup>4</sup>*Research Inst. of Organic Agriculture (FiBL), International Cooperation, Switzerland*

Compost plays a crucial role in sustainable agriculture as a rich source of organic matter, nutrients, and biota that improve soil health. Materials like *Lantana camara* and *Tithonia diversifolia* are readily available in the tropics to supplement nutrients in cattle manure during composting. There is however limited information on the succession of nutrients and microbes in compost as influenced by these materials. Moreover, prior data from the long-term farming system comparison trials in Kenya has shown significant mass and nutrient losses in compost. Compost emits GHGs which may be attributable to the nature of feedstock, duration of composting, and poor composting practices. This study sought to enumerate the contribution of composting material and duration on biological and nutrient succession in manure. A field experiment was conducted involving treatments based on grass clippings, *Lantana*, and *Tithonia* twigs with data collection every 21 days over 84 days. Other materials used in combinations were; fresh cow-dung and dry maize stalk in a ratio of 4:2:1 against the treatment material as per farmer practice. Our study showed significant losses of GHGs and nutrients during the early stages of composting with stability of these elements recorded towards the end of the composting period. However, significantly higher microbial populations were recorded at this stage but more diversity was observed at the later stages of composting. *Lantana*-based compost had the highest microbial diversity, yet the lowest abundance among all compost treatments, implying the need for specialised microbes to break down this complex material compared to other sources. This makes it suitable for the introduction of diverse microbes to degraded soils and ecosystems but unsuitable in ecosystems requiring a rapid increase in microbial populations. Overall, the study showed that composting days had an overarching influence on nutrient evolution. We recommend mitigation measures to reduce nutrient losses at the early stages of composting.

**Keywords:** Biological stability, compost quality, composting duration, GHGs, green materials, nutrient losses

---

**Contact Address:** Felix Matheri, International Centre of Insect Physiology and Ecology (icipe), Nairobi, Kenya, e-mail: fmatheri@icipe.org